

Photosynthesis

- $6 \text{ CO}_2 + 6 \text{ H}_2\text{O} + \text{light energy} \rightarrow \text{C}_6\text{H}_{12}\text{O}_6 + 6 \text{ O}_2$
- Photosynthesis is a complex biochemical pathway.
- Photosynthesis consists of two independent pathways called the **light-dependent reaction** (light reaction) and the **light-independent reaction** (dark reaction).
 - Light Reactions: the energy in sunlight is trapped, O_2 is released, and both ATP and NADPH + H^+ (hydrogen-carrier molecule) are formed
 - Dark Reactions: the ATP and NADPH + H^+ react with CO_2 from the atmosphere and form glucose
- The entire process results in the transformation of light energy from the sun into energy stored in the bonds of the glucose molecule.

Structure of a Chloroplast

- double membrane
- thylakoid: a flattened sac that contains chlorophyll
- lumen: an internal reservoir in a thylakoid
- granum: a stack of thylakoids in a chloroplast
- stroma: a solution surrounding the grana

The light reactions take place in the thylakoid membrane

The dark reactions take place in the stroma

Chlorophyll & Accessory Pigments

- Photosynthesis requires the presence of pigments that are acted upon by sunlight.
 - Sunlight consists of particles of energy that move in waves of different wavelengths (visible light is part of the electromagnetic spectrum).
 - Wavelength (λ) is the distance between the crests of successive waves.
 - The shorter the wavelength, the more energy the light has.
 - Therefore, violet light (shorter λ) has more energy than red light (longer λ).
 - Pigments are light-absorbing compounds.
 - Pigments appear colored because they absorb light of certain wavelengths and reflect that of others.
 - Chlorophyll *a* is the primary pigment in green plants that absorbs red and blue/violet light and reflects green light.
 - Chloroplasts also contain other pigments called accessory pigments.
 - Accessory pigments trap wavelengths of light that cannot be absorbed by chlorophyll *a* and then transfer the energy to chlorophyll *a* molecules for use in photosynthesis. In this way, accessory pigments enable plants to use a greater amount of the sun's energy than is available to chlorophyll alone.
 - The most common accessory pigments in green plants are chlorophyll *b* (green), carotenes (orange), xanthophylls (yellow), and anthocyanins (red)

Light Reaction (Light-Dependent Reaction)

- Pigments that are in the chloroplasts intercept light and begin the light reactions of photosynthesis.
- The light reactions occur in two photosystems (located in the thylakoid membrane):
 - Photosystem: a unit of several hundred chlorophyll *a* molecules and associated acceptor molecules

-photosystem I (**PSI**)
-photosystem II (**PSII**)

- **Steps:**

1. Sunlight strikes PSII. The light energy boosts the energy level of the electrons in the chlorophyll molecules to such a high level that the electrons can escape the chlorophyll.
 2. When light struck PSII, a water molecule split into 2H^+ , 1 O, and 2 electrons. These electrons pay back the two electrons that PSII lost in Step 1. The hydrogen ions join other hydrogen ions in the lumen. The oxygen unites with another oxygen atom and O_2 is released into the atmosphere.
 3. Two excited electrons pass down the electron transport chain (a series of electron carrier molecules) to PSI.
 4. As the electrons move along the electron transport chain, hydrogen ions (H^+) are pumped across the thylakoid membrane from the stroma. The energy that the electrons release as they flow down the ETC is used to power this H^+ pump.
 5. Sunlight strikes PSI (remember that PSI has two extra electrons from PSII). The two excited electrons pass through a series of molecules until they reach an electron acceptor called NADP^+ . Two electrons and two hydrogen ions (H^+) then attach to NADP^+ to form $\text{NADPH} + \text{H}^+$ (a product of the light reaction).
 6. The pumping of hydrogen ions into the lumen of the thylakoid creates an electrochemical gradient. As the hydrogen ions flow down the gradient from the lumen to the stroma, ADP is phosphorylated to ATP by ATP synthase (enzyme/channel). This process is called **chemiosmosis**.
- Products: ATP, NADPH + H^+ , and O_2
 - ATP and NADPH + H^+ are released into the stroma, and are used to power the dark reaction.

Dark Reaction (Light Independent Reaction)

- The dark reaction is also known as the Calvin Cycle, after an American scientist (Melvin Calvin) who figured out the pathway.
- The dark reaction is also known as the C_3 Cycle because the first stable products of this pathway are molecules that contain three carbon atoms.
- The Calvin Cycle occurs in the stroma.
- **Steps:**
 1. *CO_2 is fixed by ribulose biphosphate (RuBP)*
 CO_2 enters the plant from the atmosphere. RuBP (a five-carbon sugar molecule) binds to CO_2 . This is called CO_2 fixation. RuBP carboxylase catalyzes this reaction. The addition of CO_2 to RuBP forms an unstable six-carbon sugar molecule that immediately splits into two molecules of phosphoglyceric acid (PGA), a three-carbon molecule.
 2. *Phosphoglyceraldehyde (PGAL) is formed*
 PGAL is formed through the following 2 step process:
 -A PGA molecule binds with a phosphate group supplied by ATP.
 $\text{PGA} + \text{ATP} \rightarrow \text{PGA}\sim\text{P} + \text{ADP}$
 -Then the molecule reacts with $\text{NADPH} + \text{H}^+$, breaking the phosphate bond and forming PGAL (three-carbon molecule).
 $\text{PGA}\sim\text{P} + \text{NADPH} + \text{H}^+ \rightarrow \text{PGAL} + \text{NADP}^+ + \text{H}^+$
 3. *Glucose is formed*
 2 molecules of PGAL combine to form a molecule of glucose.
 4. *RuBP is regenerated*
 The remaining 10 molecules of PGAL are used to regenerate more RuBP, using energy supplied by ATP.
- Six turns of the Calvin Cycle are required to produce one molecule of glucose.
- Product: Glucose